Category

Synthesis of Heterocycles

Key words

enynes cobalt catalysis hydroboration γ-lactams C. WANG, S. GE* (NATIONAL UNIVERSITY OF SINGAPORE, SINGAPORE)

Versatile Cobalt-Catalyzed Enantioselective Entry to Boryl-Functionalized All-Carbon Quaternary Stereogenic Centers

J. Am. Chem. Soc. 2018, 140, 10687-10690.

Cobalt-Catalyzed Hydroboration/Cyclization Synthesis of γ-Lactams

Significance: Compounds containing the γ -lactam ring are widely distributed in nature and are present in many biologically active compounds. Besides other methods described for preparing this scaffold, transition-metal-catalyzed cyclization has gained much attention (S. Soleimani-Amiri et al. *RSC Adv.* **2017**, 7, 28407). Wang and Ge report a Co-catalyzed preparation of chiral γ -lactams bearing a boron functionality that has not previously been described and that adds to the limited number of reported reactions for the preparation of all-carbon quaternary stereogenic centers.

1,6-enynes 1 to produce γ-lactams 2. Substrates with aliphatic or aromatic substituents on the nitrogen atom and the allyl group (R^1 , R^2 = alkyl, aryl) were suitable for the reaction conditions. O- and N-tethered 1,6-enynes 1 also gave the corresponding cyclization products 2 efficiently. Steric hindrance around the nitrogen atom led to slightly decreased enantioselectivities. Enynes 1 (Ar = EWG- or EDG-bearing aryl or hetaryl) also gave products 2 in good yields and with high enantioselectivities. A gram-scale synthesis of 2 and a series of transformations including oxidations, homologations, cross-couplings, and alkenylations demonstrated the synthetic utility of this method. A reaction mechanism was proposed on the basis of deuterium-labeling and control experiments.

Comment: Reported is the asymmetric Co-cata-

lyzed hydroboration/cyclization of amide-tethered

SYNFACTS Contributors: Victor Snieckus, Lívia C. R. M. da Frota Synfacts 2018, 14(11), 1122 Published online: 18.10.2018 **DOI:** 10.1055/s-0037-1611252; **Reg-No.:** V11618SF